

Colorado **Water Supply Outlook Report**

January 1, 2019



The Sneffels Range which form the northern front of the San Juan Mountains viewed from the Uncompangre Plateau on Christmas Day 2018. While snowpack is still well below normal in the southwest corner of Colorado storms right after Christmas and over the New Year helped to improve the snowpack.

Photo By: Karl Wetlaufer Date: December 25th, 2018

REMINDER: We are soliciting field work photos from the field again this year. Each month we will pick one to grace the cover of this report! Please include information on where, when and of who/what the photo was taken.


Contents

Colorado Statewide Water Supply Conditions	3
Summary	3
Snowpack	4
Precipitation	5
Reservoir Storage	6
Streamflow	7
GUNNISON RIVER BASIN	8
UPPER COLORADO RIVER BASIN	2
SOUTH PLATTE RIVER BASIN	2
YAMPA, WHITE, NORTH PLATTE, AND LARAMIE RIVER BASINS	2
ARKANSAS RIVER BASIN	2
UPPER RIO GRANDE RIVER BASIN	2
SAN MIGUEL, DOLORES, ANIMAS, AND SAN JUAN RIVER BASINS	2
How to Read Snowpack Graphs	2
How Forecasts Are Made	1
Interpreting the Forecast Graphics	1

The U.S. Department of Agriculture (USDA) prohibits discrimination against its customers. If you believe you experienced discrimination when obtaining services from USDA, participating in a USDA program, or participating in a program that receives financial assistance from USDA, you may file a complaint with USDA. Information about how to file a discrimination complaint is available from the Office of the Assistant Secretary for Civil Rights. USDA prohibits discrimination in all its programs and activities on the basis of race, color, national origin, age, disability, and where applicable, sex (including genderidentity and expression), marital status, familial status, parental status, religion, sexual orientation, political beliefs, genetic information, reprisal, or because all or part of an individual's income is derived from any public assistance program. (Not all prohibited bases apply to all programs.) To file a complaint of discrimination, complete, sign, and mail a program discrimination complaint form, available at any USDA office location or online at www.ascr.usda.gov, or write to: USDA Office of the Assistant Secretary for Civil Rights 1400 Independence Avenue, SW. Washington, DC 20250-9410. Or call toll free at (866) 632-9992 (voice) to obtain additional information, the appropriate office or to request documents. Individuals who are deaf, hard of hearing, or have speech disabilities may contact USDA through the Federal Relay service at (800) 877-8339 or (800) 845-6136 (in Spanish). USDA is an equal opportunity provider, employer, and lender. Persons with disabilities who require alternative means for communication of program information (e.g., Braille, large print, audiotape, etc.) should contact USDA's TARGET Center at (202) 720-2600 (voice and TDD).


Colorado Statewide Water Supply Conditions

Summary




It has been an interesting start to water year 2019 so far across Colorado. While water year precipitation and snow accumulation have been dramatically better than last year on a statewide basis they are still not above average and the areas that were the driest last year are currently the most below normal. As of January 1st statewide water year precipitation was 97 percent of average. On the low end, the combined southwest basins of the San Miguel, Dolores, Animas, and San Juan have received 80 percent of normal. Conversely, the South Platte has received the highest percent of normal water year precipitation at 114 percent. Snowpack reflects similar trends across the state but at slightly lower levels than precipitation in most basins, relative to normal. Snowpack is at a high of 116 percent of normal in the Arkansas River basin and a low of 66 percent in the southwest basins. Reservoir storage also varies quite widely across the state but some large ones have been notably lacking. Blue Mesa, Colorado's largest reservoir, is experiencing one of its lowest volumes since the early 1980s and Ridgway Reservoir is the lowest since it first filled in the late 1980's. While there is still a lot of winter yet to come, it would be very beneficial for this pattern to change. This is particularly true in the southern parts of the state, where last year's extremely low stream flows led to depleted reservoir storage. A good snowpack this winter would be very beneficial to ensure ample water supply for summer 2019.

Snowpack




The 2019 water year had a promising start when an early season storm favoring the southern mountains dumped snow across the state, leaving most major river basins with above normal snowpacks. Conditions have since changed for southern Colorado where November and December have brought snowfall accumulations that are close to only half of normal amounts for the Rio Grande and combined San Miguel, Dolores, Animas, and San Juan River basins. This has led to a decline in the percent of normal snowpack for these basins, from above 200 percent of median on November 1st to 70 percent or less on January 1st. While better than last year, this is not the start many are hoping for in the parched southwest, where those river basins failed to reach snowpack levels above 60 percent of normal at any point last year. The mountains in Colorado's other river basins are holding more plentiful snowpacks thanks to consistent snowfall through October and November. December accumulations were below normal across the state, but prior snowfall buffered against any major decreases in the percent of normal for most basins. The Arkansas and South Platte River basins have the most abundant snowpacks, with respect to normal, at 116 and 112 percent, respectively. The Colorado and combined Yampa, White, and North Platte River basins are also above normal at 107 and 104 percent of median, respectively, while the Gunnison River basin is maintaining a modest snowpack at 91 percent of median. Statewide, Colorado's mountain snowpack is at 93 percent of normal. About three months remain in the accumulation season before most mountain locations reach peak snowpack levels, leaving time for an uptick in snowfall to improve conditions in the southwest. About 120 percent of normal future snow accumulations are needed to bring the southern basins to normal levels, which is certainly possible. However, another month with subpar snowfall will make normal snowpack levels prior to runoff difficult to achieve.

Precipitation



Water Year 2019 got off to a great start with all major basins receiving above average precipitation in October. This ranged from a low of 109 percent of average in the combined Yampa, White, and North Platte basins to a high of 144 percent in the combined San Miguel, Dolores, Animas, and San Juan basins of Southwest Colorado. November precipitation displayed notable differences between the northern and southern parts of the state. Northern Colorado continued to receive well above average precipitation while the Rio Grande and combined southwest basins got a meager 53 and 40 percent of average precipitation, respectively. Precipitation patterns shifted again in December with all of Colorado receiving below average precipitation in a more consistent manner. All major basins were in the range of 57 to 74 percent of average precipitation for the month. Statewide this has left Colorado with near normal water-year-to-date precipitation, at 97 percent of average. Largely driven by the differences observed in November, central and northern Colorado generally has near to above average water year precipitation and most of southern Colorado has lower than average water year precipitation. The South Platte and Upper Colorado basins have the highest values at 114 and 107 percent of normal, respectively. The Combined Yampa, White, and North Platte along with the Arkansas and Gunnison basins are near normal with 101, 98, and 97 percent of average, respectively. That leaves the Upper Rio Grande and basins of Southwest Colorado with the least water year precipitation, at 84 and 80 percent, respectively.

Reservoir Storage




While reservoir storage is near to well below average in all major basins of Colorado, every basin except the Arkansas did show some degree of increase in storage over the last month, relative to normal values. Currently statewide reservoir storage is 81 percent of average. The South Platte and combined Yampa and White basins have close to normal storage for this time of year at 99 and 97 percent of average, respectively. The Arkansas and Colorado basins are holding slightly lower values but still relatively close to normal, at 92 and 91 percent of average, respectively. The Upper Rio Grande basin is in the middle of the pack, holding 80 percent of average storage. While this basin doesn't have the most total storage they are at least in a much better position than other basins that also had a notably dry water year 2018. The Gunnison Basin is holding only 58 percent of average storage which is 39 percent of capacity. A large driver of this is from Blue Mesa Reservoir, the largest in the state, which has been at the lowest levels since 1984 over the last several months. Storage in the combined San Miguel, Dolores, Animas, and San Juan basins is currently even slightly lower, at 56 percent of average. All four of these major rivers observed their third lowest April-July streamflows last summer with periods of record ranging from 61 to 107 years leading to sharp declines in reservoir storage. Given these very low values across Southwest Colorado another year of low snowpack could lead to challenging water resource issues next summer.

Streamflow

Due to current staffing, most official forecasts only will be available February through May. If you rely on the January or June forecasts, please contact cara.s.mccarthy@por.usda.gov or Brian.Domonkos@co.usda.gov.


In the river basin sections of the report that follow, the typical graphic displaying the range of forecast exceedance probabilities has been replaced with a new chart. The new graphic displays the historic range of observed streamflows for all the forecast points in each of the major river basins. In the example image provided below, the red line represents the minimum observed volume; the green line, average volume; and the blue line, the maximum observed streamflow volume. The numbers below the lines are the observed streamflow volumes in thousand acre-feet (KAF) followed by the year of observed flow in parentheses. At the end of this report, there is further explanation of these graphics for periods when streamflow forecasts are available.




GUNNISON RIVER BASIN


January 1, 2019

Snowpack in the Gunnison River basin is below normal at 91% of the median. Precipitation for December was 74% of average which brings water year-to-date precipitation to 97% of average. Reservoir storage at the end of December was 58% of average compared to 104% last year. No forecasts are available for January.



^{*}SWE values calculated using daily SNOTEL data only






Watershed Snowpack Analysis January 1st, 2019

Last Year %

Sub-Basin	# of Sites	% Median	Median
Upper Gunnison	10	94	41
Surface Creek	2	99	16
Uncompahgre	3	82	26
Basin-Wide Total	13	91	37


^{*}SWE values calculated using first of month SNOTEL data and snow course measurements



Reservoir Storage End of December 2018

	Current	Last Year	Average	Capacity
Reservoir	(KAF)	(KAF)	(KAF)	(KAF)
BLUE MESA RESERVOIR	248.8	592.6	549.9	830.0
CRAWFORD RESERVOIR	1.2	4.6	7.1	14.0
CRYSTAL RESERVOIR	7.1	8.2	7.7	17.5
FRUITGROWERS RESERVOIR	1.7	1.6	2.8	3.6
FRUITLAND RESERVOIR	0.3	0.5	1.0	9.2
MORROW POINT RESERVOIR	107.6	111.0	111.6	121.0
PAONIA RESERVOIR	2.6	2.1	3.5	15.4
RIDGEWAY RESERVOIR	45.9	60.3	68.8	83.0
SILVERJACK RESERVOIR	0.8	2.2	5.0	12.8
TAYLOR PARK RESERVOIR	59.0	76.2	68.1	106.0
VOUGA RESERVOIR	0.2	0.6	0.7	0.9
BASINWIDE	475.1	859.8	826.2	1213.4
Number of Reservoirs	11	11	11	11


Due to current staffing, most official forecasts only will be available February through May. If you rely on the January or June forecasts, please contact cara.s.mccarthy@por.usda.gov or Brian.Domonkos@co.usda.gov. The following graphic displays the historic range of observed streamflows for reference during months where forecasts are not available. Red = minimum, green = average, and blue = maximum observed flow.




UPPER COLORADO RIVER BASIN


January 1, 2019

Snowpack in the Colorado River basin is above normal at 107% of the median. Precipitation for December was 69% of average which brings water year-to-date precipitation to 107% of average. Reservoir storage at the end of December was 91% of average compared to 113% last year. No forecasts are available for January.



^{*}SWE values calculated using daily SNOTEL data only






Watershed Snowpack Analysis January 1st, 2019

Last Year %

Sub-Basin	# of Sites	% Median	Median
Blue River	5	124	87
Upper Colorado	19	107	78
Muddy Creek	3	109	96
Eagle River	4	98	48
Plateau Creek	5	94	27
Roaring Fork	7	107	60
Williams Fork	3	104	64
Willow Creek	2	113	119
Basin-Wide Total	28	107	69


^{*}SWE values calculated using first of month SNOTEL data and snow course measurements



Reservoir Storage End of December 2018

	Current	Last Year	Average	Capacity
Reservoir	(KAF)	(KAF)	(KAF)	(KAF)
DILLON RESERVOIR	177.7	236.3	222.1	249.1
LAKE GRANBY	348.6	437.4	325.7	465.6
GREEN MOUNTAIN RESERVOIR	51.0	66.7	85.2	146.8
HOMESTAKE RESERVOIR	41.2	41.1	31.9	43.0
RUEDI RESERVOIR	60.0	72.1	76.8	102.0
VEGA RESERVOIR	5.0	9.6	11.8	32.9
WILLIAMS FORK RESERVOIR	69.7	66.0	66.5	97.0
WILLOW CREEK RESERVOIR	6.4	5.9	6.6	9.1
WOLFORD MOUNTAIN RESERVOIR	34.7	55.0	44.0	65.9
SHADOW MOUNTAIN RESERVOIR	17.3	17.3	17.3	18.4
BASINWIDE	811.7	1007.4	887.9	1229.8
Number of Reservoirs	10	10	10	10


Due to current staffing, most official forecasts only will be available February through May. If you rely on the January or June forecasts, please contact cara.s.mccarthy@por.usda.gov or Brian.Domonkos@co.usda.gov. The following displays the historic range of observed streamflows for reference during months where forecasts are not available.




SOUTH PLATTE RIVER BASIN

January 1, 2019

Snowpack in the South Platte River basin is above normal at 112% of the median. Precipitation for December was 62% of average which brings water year-to-date precipitation to 114%. Reservoir storage at the end of December was 99% of average compared to 114% last year. No forecasts are available for January.



*SWE values calculated using daily SNOTEL data only **Mountain Precipitation** 160 140 120 Percent of Average 100 80 60 40 20 0 Oct Nov Dec Jan Feb Mar Apr May Monthly Year-to-date




Watershed Snowpack Analysis January 1st, 2019

Last Year %

Sub-Basin	# of Sites	% Media	an	Median	
Big Thompson	:	3	106		80
Boulder Creek	:	3	104		92
Cache La Poudre	:	2	100		100
Clear Creek	:	2	107		86
Saint Vrain		1	133		67
Upper South Platte	(5	132		82
Basin-Wide Total	1	7	112		86

^{*}SWE values calculated using first of month SNOTEL data and snow course measurements



Average

Capacity


Reservoir Storage End of December 2018

Current

Last Year

	Current	Last Year	Average	Capacity
Reservoir	(KAF)	(KAF)	(KAF)	(KAF)
ANTERO RESERVOIR	18.9	20.1	15.5	19.9
BARR LAKE	16.8	26.9	22.3	30.1
BLACK HOLLOW RESERVOIR	4.1	3.4	2.8	6.5
BOYD LAKE	31.5	33.0	27.4	48.4
CACHE LA POUDRE	6.1	8.9	5.4	10.1
CARTER LAKE	58.7	49.3	67.5	108.9
CHAMBERS LAKE	3.3	6.4	3.1	8.8
CHEESMAN LAKE	60.0	72.2	64.3	79.0
COBB LAKE	15.2	19.2	11.7	22.3
ELEVENMILE CANYON RESERVOIR	99.2	100.0	95.9	98.0
EMPIRE RESERVOIR	24.0	25.1	20.6	36.5
FOSSIL CREEK RESERVOIR	9.2	9.3	6.3	11.1
GROSS RESERVOIR	14.9	18.6	15.4	29.8
HALLIGAN RESERVOIR	4.1	6.4	3.9	6.4
HORSECREEK RESERVOIR	0.0	7.7	8.5	14.7
HORSETOOTH RESERVOIR	74.0	76.5	83.5	149.7
JACKSON LAKE RESERVOIR	24.3	24.2	20.9	26.1
JULESBURG RESERVOIR	16.5	16.5	17.0	20.5
LAKE LOVELAND RESERVOIR	0.0	6.3	6.8	10.3
LONE TREE RESERVOIR	6.5	6.9	5.7	8.7
MARIANO RESERVOIR	0.7	4.0	2.9	5.4
MARSHALL RESERVOIR	5.2	6.6	5.4	10.0
MARSTON RESERVOIR	10.3	7.2	6.0	13.0
MILTON RESERVOIR	18.7	19.1	14.3	23.5
POINT OF ROCKS RESERVOIR	47.2	51.7	43.3	70.6
PREWITT RESERVOIR	16.3	21.1	13.9	28.2
RIVERSIDE RESERVOIR	41.2	47.0	32.1	55.8
SPINNEY MOUNTAIN RESERVOIR	27.0	36.3	30.5	49.0
STANDLEY RESERVOIR	29.0	41.2	35.8	42.0
TERRY RESERVOIR	4.8	5.8	5.1	8.0
UNION RESERVOIR	8.9	11.1	9.8	13.0
WINDSOR RESERVOIR	7.9	9.8	7.7	15.2
BASINWIDE	704.5	797.7	711.3	1079.5
Number of Reservoirs	32	32	32	32


Due to current staffing, most official forecasts only will be available February through May. If you rely on the January or June forecasts, please contact cara.s.mccarthy@por.usda.gov or Brian.Domonkos@co.usda.gov. The following displays the historic range of observed streamflows for reference during months where forecasts are not available.





YAMPA, WHITE, NORTH PLATTE, AND LARAMIE RIVER BASINS


January 1, 2019

Snowpack in the Yampa, White & North Platte basins is above normal at 104% of the median. Precipitation for December was 64% of average and water year-to-date precipitation is 101% of average. Reservoir storage at the end of December was 97% of average compared to 117% last year. No forecasts are available for January.










Watershed Snowpack Analysis January 1st, 2019

Last Year %

Sub-Basin	# of Sites	% Median	Median
Laramie	2	94	118
North Platte	8	106	83
Total Laramie & North Platte	10	104	89
Elk	2	90	68
Yampa	9	107	73
White	3	107	60
Total Yampa & White	11	105	68
Little Snake	7	103	71
Basin-Wide Total	25	104	77


^{*}SWE values calculated using first of month SNOTEL data and snow course measurements



Reservoir Storage End of December 2018

	Current	Last Year	Average	Capacity
Reservoir	(KAF)	(KAF)	(KAF)	(KAF)
STAGECOACH RESERVOIR NR OAK C	31.2	33.4	29.3	36.5
YAMCOLO RESERVOIR	2.5	7.1	5.3	8.7
BASINWIDE	33.7	40.5	34.6	45.2
Number of Reservoirs	2	2	2	2


Due to current staffing, most official forecasts only will be available February through May. If you rely on the January or June forecasts, please contact cara.s.mccarthy@por.usda.gov or Brian.Domonkos@co.usda.gov. The following displays the historic range of observed streamflows for reference during months where forecasts are not available.




ARKANSAS RIVER BASIN


January 1, 2019

Snowpack in the Arkansas River basin is above normal at 116% of the median. Precipitation for December was 61% of average which brings water year-to-date precipitation to 98% of average. Reservoir storage at the end of December was 92% of average compared to 139% last year. No forecasts are available for January.



*SWE values calculated using daily SNOTEL data only






Watershed Snowpack Analysis January 1st, 2019

Last Year %

				Lastical	/0
Sub-Basin	# of Sites	% Med	ian	Median	
Upper Arkansas		3	121		81
Cucharas & Huerfano		3	111	-	17
Purgatoire		2	171	-	21
Basin-Wide Total		8	116	i	48


^{*}SWE values calculated using first of month SNOTEL data and snow course measurements



Reservoir Storage End of December 2018

	Current	Last Year	Average	Capacity
Reservoir	(KAF)	(KAF)	(KAF)	(KAF)
ADOBE CREEK RESERVOIR	9.0	47.7	32.7	62.0
CLEAR CREEK RESERVOIR	6.6	7.1	6.7	11.4
CUCHARAS RESERVOIR				40.0
GREAT PLAINS RESERVOIR				150.0
HOLBROOK LAKE	0.3	6.2	2.5	7.0
HORSE CREEK RESERVOIR	13.6	26.9	9.4	27.0
JOHN MARTIN RESERVOIR	146.2	268.9	122.8	616.0
LAKE HENRY	4.8	9.0	3.7	9.4
MEREDITH RESERVOIR	21.7	39.1	19.7	42.0
PUEBLO RESERVOIR	196.8	220.6	170.8	354.0
TRINIDAD LAKE	20.5	38.3	24.4	167.0
TURQUOISE LAKE	73.2	93.4	94.1	127.0
TWIN LAKES RESERVOIR	36.3	42.2	57.0	86.0
BASINWIDE	529.0	799.4	543.8	1698.8
Number of Reservoirs	11	11	11	13


Due to current staffing, most official forecasts only will be available February through May. If you rely on the January or June forecasts, please contact cara.s.mccarthy@por.usda.gov or Brian.Domonkos@co.usda.gov. The following displays the historic range of observed streamflows for reference during months where forecasts are not available.




UPPER RIO GRANDE RIVER BASIN


January 1, 2019

Snowpack in the Upper Rio Grande River basin is below normal at 70% of median. Precipitation for December was 58% of average which brings water year-to-date precipitation to 84% of average. Reservoir storage at the end of December was 80% of average compared to 123% last year. No forecasts are available for January.



^{*}SWE values calculated using daily SNOTEL data only






Watershed Snowpack Analysis January 1st, 2019

Last Year %

Sub-Basin	# of Sites	% Median	Median
Alamosa Creek	1	44	52
Conejos & Rio San Antonio	2	53	42
Culebra & Trinchera Creek	3	95	26
Upper Rio Grande	6	64	26
Basin-Wide Total	12	70	29


^{*}SWE values calculated using first of month SNOTEL data and snow course measurements



Reservoir Storage End of December 2018

	Current	Last Year	Average	Capacity
Reservoir	(KAF)	(KAF)	(KAF)	(KAF)
CONTINENTAL RESERVOIR	14.6	10.8	3.8	27.0
PLATORO RESERVOIR	19.3	23.3	24.0	60.0
RIO GRANDE RESERVOIR	0.0	25.8	14.8	51.0
SANCHEZ RESERVOIR	7.0	21.2	27.5	103.0
SANTA MARIA RESERVOIR	24.0	18.9	10.4	45.0
TERRACE RESERVOIR	3.4	7.5	5.5	18.0
BEAVER RESERVOIR	3.6	3.3	4.1	4.5
BASINWIDE	72.0	110.7	90.1	308.5
Number of Reservoirs	7	7	7	7


Due to current staffing, most official forecasts only will be available February through May. If you rely on the January or June forecasts, please contact cara.s.mccarthy@por.usda.gov or Brian.Domonkos@co.usda.gov. The following displays the historic range of observed streamflows for reference during months where forecasts are not available.




SAN MIGUEL, DOLORES, ANIMAS, AND SAN JUAN RIVER BASINS


January 1, 2019

Snowpack in the combined southwest river basins is below normal at 66% of median. Precipitation for December was 57% of average which brings water year-to-date precipitation to 80% of average. Reservoir storage at the end of December was 56% of average compared to 105% last year. No forecasts are available for January.



^{*}SWE values calculated using daily SNOTEL data only






Watershed Snowpack Analysis January 1st, 2019

Last Year %

Sub-Basin	# of Sites	% Median	Median
Animas	9	74	24
Dolores	5	59	19
San Miguel	3	66	18
San Juan	3	55	25
Basin-Wide Total	19	66	22

^{*}SWE values calculated using first of month SNOTEL data and snow course measurements



Reservoir Storage End of December 2018

	Current	Last Year	Average	Capacity
Reservoir	(KAF)	(KAF)	(KAF)	(KAF)
GROUNDHOG RESERVOIR	0.2	12.1	12.3	22.0
JACKSON GULCH RESERVOIR	1.7	5.2	4.5	10.0
LEMON RESERVOIR	6.9	18.7	20.7	40.0
MCPHEE RESERVOIR	168.0	285.5	265.6	381.0
NARRAGUINNEP RESERVOIR	2.0	11.7	14.1	19.0
VALLECITO RESERVOIR	33.8	64.9	62.4	126.0
TROUT LAKE RESERVOIR	2.1	2.6	2.5	3.2
BASINWIDE	214.7	400.8	382.1	601.2
Number of Reservoirs	7	7	7	7

Due to current staffing, most official forecasts only will be available February through May. If you rely on the January or June forecasts, please contact cara.s.mccarthy@por.usda.gov or Brian.Domonkos@co.usda.gov. The following displays the historic range of observed streamflows for reference during months where forecasts are not available.



How to Read Snowpack Graphs

The graphs show snow water equivalent (SWE) (in inches), using daily SNOTEL data. for the October 1 through September 30 water year. Basin "observed" SWE values are computed using SNOTEL sites which are characteristic of the snowpack of the particular basin. The SWE observations at these sites are averaged and normalized to produce these basin snowpack graphs.


Current water year is represented by the heavy red line terminating on the last day the graphic was updated.

Historical observed percentile range is shown as a gray background area on the graph. Shades of gray indicate maximum, 90 percentile, 70 percentile, 50 percentile (solid black line), 30 percentile, 10 percentile, and minimum for the period of record.

50 % Excedance Projection: The most probabilistic snowpack projection, based on the median snowpack is projected forward from the end of the current period to the end of the current water year.

For more detailed information on these graphs visit:

http://www.nrcs.usda.gov/Internet/FSE DOCUMENTS/nrcs144p2 062291.pdf



How Forecasts Are Made

For more water supply and resource management information, contact:

Brian Domonkos Snow Survey Supervisor USDA, Natural Resources Conservation Service Denver Federal Center, Bldg 56, Rm 2604 PO Box 25426 Denver, CO 80225-0426 Phone (720) 544-2852

Website: http://www.nrcs.usda.gov/wps/portal/nrcs/main/co/snow/


Most of the annual streamflow in the western United States originates as snowfall that has accumulated in the mountains during the winter and early spring. As the snowpack accumulates, hydrologists estimate the runoff that will occur when it melts. Measurements of snow water equivalent at selected manual snow courses and automated SNOTEL sites, along with precipitation, antecedent streamflow, and indices of the El Niño / Southern Oscillation are used in computerized statistical and simulation models to prepare runoff forecasts. Unless otherwise specified, all forecasts are for flows that would occur naturally without any upstream influences.

Forecasts of any kind, of course, are not perfect. Streamflow forecast uncertainty arises from three primary sources: (1) uncertain knowledge of future weather conditions, (2) uncertainty in the forecasting procedure, and (3) errors in the data. The forecast, therefore, must be interpreted not as a single value but rather as a range of values with specific probabilities of occurrence. The middle of the range is expressed by the 50% exceedance probability forecast, for which there is a 50% chance that the actual flow will be above, and a 50% chance that the actual flow will be below, this value. To describe the expected range around this 50% value, four other forecasts are provided, two smaller values (90% and 70% exceedance probability) and two larger values (30%, and 10% exceedance probability). For example, there is a 90% chance that the actual flow will be more than the 90% exceedance probability forecast. The others can be interpreted similarly.

The wider the spread among these values, the more uncertain the forecast. As the season progresses, forecasts become more accurate, primarily because a greater portion of the future weather conditions become known; this is reflected by a narrowing of the range around the 50% exceedance probability forecast. Users should take this uncertainty into consideration when making operational decisions by selecting forecasts corresponding to the level of risk they are willing to assume about the amount of water to be expected. If users anticipate receiving a lesser supply of water, or if they wish to increase their chances of having an adequate supply of water for their operations, they may want to base their decisions on the 90% or 70% exceedance probability forecasts, or something in between. On the other hand, if users are concerned about receiving too much water (for example, threat of flooding), they may want to base their decisions on the 30% or 10% exceedance probability forecasts, or something in between. Regardless of the forecast value users choose for operations, they should be prepared to deal with either more or less water. (Users should remember that even if the 90% exceedance probability forecast is used, there is still a 10% chance of receiving less than this amount.) By using the exceedance probability information, users can easily determine the chances of receiving more or less wate

Interpreting the Forecast Graphics

These graphics provide a new way to visualize the range of streamflows represented by the forecast exceedance probabilities for each forecast period. The colors in the bar for each forecast point indicate the exceedance probability of the forecasts and the vertical lines on the bar signify the five published forecast exceedance probabilities. The numbers displayed above the color scale represent the actual forecasted streamflow volume (in KAF) for the given exceedance probability. The horizontal axis provides the percent of average represented by each forecast and the gray line centered above 100% represents the 1981-2010 historical average streamflow. The position of the gray line relative to the color scale provides a benchmark for considering future streamflows. If the majority of the forecast range is to the right of the gray line, there is a higher likelihood of above average streamflow volumes during the provided forecast period. Conversely, if the majority of the color bar is to the left of the average mark, below average volumes are more likely. The horizontal span of the forecasts offers an indication of the uncertainty in a given forecast: when the bar spans a large horizontal range, the forecast skill is low and uncertainty is high; when the bar is narrow in width, the forecast skill is higher and uncertainty lower.





Denver Federal Center, Bldg 56, Rm 2604 PO Box 25426 Denver, CO 80225-0426

In addition to the water supply outlook reports, water supply forecast information for the Western United States is available from the Natural Resources Conservation Service and the National Weather Service monthly, January through June. The information may be obtained from the Natural Resources Conservation Service web page at http://www.wcc.nrcs.usda.gov/wsf/westwide.html

Issued by

Matthew J. Lohr
Chief, Natural Resources Conservation Service
Farm Production and Conservation Mission Area

U.S. Department of Agriculture

Released by

Clint Evans

State Conservationist

Natural Resources Conservation Service

Lakewood, Colorado

Colorado Water Supply Outlook Report

Natural Resources Conservation Service Lakewood, CO